



Monitoring for overfeeding


1. Respiratory & Cardiac Parameters



2. Biochemistry



3. Kcal Delivery >110%



High risk patients:

- Vulnerable, elderly, frail
- Malnourished
- Acute phase

- Multiple nutrition routes
- PN
- Transitioning PN to EN
- Non-nutrition kcals

- Catch up feeding
- Everyone!

Monitoring for overfeeding

Overfeeding is harmful in critically ill patients. It is defined as providing >110% of estimated kcal requirement.

Signs of overfeeding:

1. Respiratory and cardiac parameters:

- Difficulties in respiratory weaning raised/rising pCO₂ levels, increasing minute ventilation (V_e) and FiO₂
- New pulmonary oedema
- New arrhythmias

May be difficult to determine whether occurrence of these symptoms is due to overfeeding or to COVID 19.

2. Biochemical parameters:

- Hyperglycaemia
- Raised/rising LFTs (EN and PN) – may be due to reasons other than overfeeding e.g. sepsis, antibiotic therapy, other medications, acute liver injury
- Raised triglycerides (EN and PN) – consider propofol rate also
- Refeeding dyselectrolytaemia

3. Kcal delivery

- Providing > 110% of kcal requirements
- Include non-nutritional kcal sources such as propofol, dextrose, citrate

Monitoring for underfeeding

Cumulative underfeeding is harmful to the critically ill. Record % kcal and protein delivery versus goal each review

Signs of underfeeding:

- Wt loss, appearance of wt loss, muscle/fat store depletion
- Deteriorating skin condition, pressure ulcers, wounds or drain/line sites
- Continuously not achieving estimated requirements
- Multiple feed interruptions